

Stamford Quadrangle Bedrock Geology Map w/ Explanation and Cross-Sections

Explanation
Map
Cross-Sections

NOTICE !

This document contains unpublished maps, cross- sections, and related information archived by the State Geological and Natural History Survey of Connecticut as part of the Survey Library Collection.

These materials have not been reviewed for accuracy, consistency, or completeness. For many geographic areas, more current information exists, either in published or unpublished form.

For the most part, these materials were developed under research and mapping agreements between the State Geological Survey and individual scientists, academic institutions, or graduate students. Some of these materials have been received by the State Geological Survey as donations. The veracity of the information contained within these documents is the responsibility of the authorship. The State Geological and Natural History Survey of Connecticut, does not promote or endorse this content, nor does the State Survey attest as to its level of accuracy.

These materials have been preserved under a cooperative agreement between the State Geological Survey and the US Geological Survey as part of the National Geological and Geophysical Data Preservation Program. www.datapreservation.usgs.gov


These materials are offered in the spirit of open government. Reproduction of these manuscripts was conducted to the highest practical degree, within the parameters of the funding mechanism. Original documents are available for inspection by contacting the [Connecticut State Geologist](#).

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

STATE OF CONNECTICUT
HIGHWAY DEPARTMENT

STAMFORD QUADRANGLE
CONNECTICUT—FAIRFIELD CO.
7.5 MINUTE SERIES (TOPOGRAPHIC)

NORWALK NORTH



Mapped, edited, and published by the Geological Survey

Control by USGS, USC&GS, USCE, and Connecticut Geodetic Survey

Topography by photogrammetric methods from aerial photographs

taken 1949. Field checked 1951. Revised 1960

Selected hydrographic data compiled from USC&GS Charts 221 and 222

(1959). This information is not intended for navigational purposes

Polyconic projection. 1927 North American datum

10,000-foot grid based on Connecticut coordinate system

1000-meter Universal Transverse Mercator grid ticks,

zone 18, shown in blue

Fine red dashed lines indicate selected fence and field lines where

generally visible on aerial photographs. This information is unchecked

Red tint indicates areas in which only landmark buildings are shown

UTM GRID AND 1980 MAGNETIC NORTH DECLINATION AT CENTER OF SHEET

12° 213 MILS 0.5° 17 MILS

CONTOUR INTERVAL 10 FEET
DATUM IS MEAN SEA LEVEL
DEPTH CURVES AND SOUNDS IN FEET—DATUM IS MEAN LOW WATER
SHORELINE SHOWN REPRESENTS THE APPROXIMATE LINE OF MEAN HIGH WATER
THE MEAN RANGE OF TIDE IS APPROXIMATELY 7.2 FEET

THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS
FOR SALE BY U. S. GEOLOGICAL SURVEY, WASHINGTON 25, D. C.
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

DEPT. OF GEOLOGY
WESLEYAN UNIVERSITY

Received
from
July 26, 1971



STAMFORD, CONN.
N4100—W7330/7.5


1960

AMS 6266 II SE—SERIES V816

Connecticut Geological and Natural History Survey Library
Department of Environmental Protection
79 Elm Street
Hartford, CT 06106-5127
(860) 424-3540

Schematic of cross-sections








Siscowit granite

Reddish-colored granite containing thin biotite-muscovite seams; Locally contains numerous Amphibolite pods



Dingletown Gneiss

Light gray, well foliated garnet-muscovite-biotite adamellite gneiss




Harrison gneiss
Unit 1

Dark gray biotite-hornblende-quartz-feldspar schist and gneiss; The gneiss commonly contains Oligoclase megacrysts




Harrison schist+granulite
UNIT 2

Funky-weathering garnet-muscovite-biotite-feldspar-quartz schist



Harrison Adamellite
UNIT 3

Biotite-hornblende-quartz-feldspar gneiss interlayered on various scales with well foliated garnet-muscovite-biotite adamellite gneiss containing prominent sillimanite-muscovite-quartz nodules.



South End Unit

medium-grained biotite adamellite gneiss with potassium feldspar megacrysts. Amphibolites and biotite-hornblende-quartz-feldspar schists occur locally.



Hartland Schist +
Granulite

Rusty to funky weathering garnet-biotite-muscovite-feldspar quartz schist & biotite-muscovite-quartz gneiss. Schist locally contains sillimanite



Hartland white gneiss (after Leo Wall)
white or light grey biotite-muscovite gneiss with local garnet.